


1 Degree Sequences

The *degree sequence* of a graph is the sequence of the degrees of the vertices, arranged in descending order, with repetitions as needed. For example, the degree sequence of the following graph is $(3, 2, 2, 2, 1)$.



For each of the parts below, determine if there exists a simple undirected graph G (i.e. a graph without self-loops and multiple-edges) having the given degree sequence. Justify your claim.

- (a) $(3, 3, 2, 2)$
- (b) $(3, 2, 2, 2, 2, 1, 1)$
- (c) $(6, 2, 2, 2)$
- (d) $(4, 4, 3, 2, 1)$

2 Bipartite Graph

A bipartite graph consists of 2 disjoint sets of vertices (say L and R), such that no 2 vertices in the same set have an edge between them. For example, here is a bipartite graph (with $L = \{\text{green vertices}\}$ and $R = \{\text{red vertices}\}$), and a non-bipartite graph.

Prove that a graph has no tours of odd length if it is a bipartite (This is equivalent to proving that, a graph G being a bipartite implies that G has no tours of odd length).

3 Not everything is normal: Odd-Degree Vertices

Claim: Let $G = (V, E)$ be an undirected graph. The number of vertices of G that have odd degree is even.
Prove the claim above using:

- (i) Direct proof (e.g., counting the number of edges in G). *Hint: in lecture, we proved that $\sum_{v \in V} \deg v = 2|E|$.*

- (ii) Induction on $m = |E|$ (number of edges)



Figure 1: A bipartite graph (left) and a non-bipartite graph (right).

(iii) Induction on $n = |V|$ (number of vertices)